

2015

1. Comparison Of dexmedetomidine 5mg Clonidine 75 Mcg added to 12.5mg Of 0.5% Bupivacaine For Spinal Anaesthesia In Lower Abdominal Surgeries. IJCR: 2015OCT: VOL 7 : ISSUE 10:PP 21318-21321.
2. Effect Of Pre Operative I.VParacetamol On Post Operative Analgesia Requirement In Patients Undergoing Laparoscopic Procedures. IJSS 2015 NO: VO13: ISSUES: P92-96.
3. A comparative study of trace element levels in coronary artery tissue of coronary heart disease patient with serum levels in healthy individuals . International Archives of integrated Medicine, vol -2 ,issue-5,may -2015.
4. Comparison of Dexmedetomidine and Clonidine ($\alpha 2$ agonist drugs) in Laparoscopic Appendectomy Under G. A. Journal of Evidence based Medicine and Healthcare; Vol 2, Issue 30, July 27, 2015; Page: 4419-4427.

2016

1. Low dose fentanyl attenuates hypertension but not tachycardia during laryngoscopy and tracheal intubation in a three arm study. IAIM, 2016; 3(4): 72-77. Page 72.
2. Efficacy of IV paracetamol as post operative analgesia for thyroid surgeries under general anaesthesia. IJSR,FEB;2016,VOL;5,ISSUE;2
3. A study to clinically evaluate the efficacy and safety of dexmedetomidine(1 μ g/kg) when used as an adjuvant to caudal ropivacaine(0.25%1 ml/kg) for infraumbilical surgeries in paediatric age group". Ijca Vol.4, Issue. 4
4. A comparative study of ultrasound guided supraclavicular brachial plexus block using bupivacaine-lignocaine with adrenaline and dexmedetomidine added to bupivacaine-lignocaine with adrenaline. IJCR;2016;8(7):69-73

2017

1. A Study to Observe Effect Of Mgso4 to 0.5% Bupivaccaine For Intrathecal Anaesthesia For Lower Limb Surgeries.IJCA. VOL. 4 : ISSUE 3 (JULY-SEPTEMBER),2017.
2. A Comparative Study Between Intrathecal Isobaric Ropivacaine 0.75% Plus Dexmedetomidine And Isobaric Ropivaccaine 0.75%Plus Fentanyl For Lower Limb Surgeries .AER: JULY-SEPTEMBER ISSUE,2017.
3. A Comparative Study BetweenRopivaccaine With Dexmeditomidine And Ropivaccaine With Fentanyl in Lower Abdominal And Lower Limb Surgeries For Post Operative Epidural Analgesia .IJAA: VOL. 4. NO.2. APR –JUNE 2017.
4. A comparative study between 0.375% ropivacaine with 50mcg of dexmedetomidine and 0.375% ropivacaine with 8mg of dexamethasone for ultrasound guided supraclavicular brachial plexus block. IJCA;2017;4(1);69—73.
5. A Comparative Study between Oral Pregabalin 150mg and Gabapentin 300mg for Postoperative Analgesia in Patients Undergoing laproscopic cholecystectomy. ijaa.2349.8471.42(pt-II)17.27.
6. USG guided femoral nerve block for fracture femur to make the patient sit comfortably for spinal Anaesthesia. Indian Journal of Anaesthesia and analgesia. 2017, 4(3 part -1),664-668.
7. right ventricular systolic function in off pump coronary artery bypass (OPCAB) and Its Measurement By Tricuspid Annular plane systolic Excursion. Indian journal of Anaesthsia and Analgesia volume - 4,Number-2,April-June 2017 (part-2).